



# AROMABIOTIC POUL 60

## Analysis

ME poultry	MJ/kg	22.3
ME broilers	MJ/kg	21.5
Humidity	g/kg	196.0
Crude protein	g/kg	4.2
Crude fat	g/kg	540
Starch	g/kg	47
Sugars + Starch	g/kg	47
Crude ash	g/kg	334
Crude fiber	g/kg	0.5
Pd p	g/kg	0.01
Calcium	g/kg	0.02
Phosphor	g/kg	0.06
Pd m	g/kg	0.01
Magnesium	g/kg	0.02
Sodium	g/kg	0.01
NaCl	g/kg	0.01
Cl	g/kg	0.01
Potassium	g/kg	0.08
Linoleic acid	g/kg	0.08
Lysin	g/kg	0.20
Methionin	g/kg	0.09
Cystin	g/kg	0.09
Met + Cys	g/kg	0.18
Threonin	g/kg	0.16
Tryptophane	g/kg	0.05
Partikal size	> 0.25mm	68.99%
	>0.5 mm	2.825%
	en < 1 mm	
	>1.00mm	0.109%

## COMPLEMENTARY FEED FOR POULTRY, FISH AND CRUSTACEA

### USE

0.4-2.5 kg per Mt final feed  
depending on the age of the  
animal

Heat stable at expansion,  
extrusion and pelleting

### TRANSPORT AND CONDITIONS OF DELIVERY

25 kg

### STORAGE AND PRESERVATION CONDITIONS

Store cool and dry

### ARTICLE NUMBER

80970001

80970000

### DATE

28/04/2009

Complies with KB of 8 February 1999 (BS 21.04.99) and MB of 12 February 1999 (BS 21.04.99) concerning trade and use of substances destined for animal feeding.